MORTALITY FROM SMOKING IN DEVELOPED COUNTRIES 1950–2000

(2nd edition, revised June 2006: www.deathsfromsmoking.net)

Richard Peto, Alan D Lopez,
Jillian Boreham and Michael Thun

Trends in annual nos. of deaths: smoking-attributed and total

One pair of pages for each of the following:

All Developed Countries Czech Republic Latvia **Russian Federation EU15*** (European Union) Denmark Lithuania Slovakia EU10[†] (European Union) **Spain** Estonia Luxembourg EU25[‡] (European Union) **Finland** Malta Sweden **Australia** Moldova **Switzerland** France Austria **Netherlands** Ukraine Germany **Belarus** Greece **New Zealand United Kingdom Belgium Norway United States** Hungary Bulgaria Ireland Poland Canada Italy **Portugal Central Asia** Japan Romania

^{*15} countries: Austria, Belgium, Denmark, Finland, France, Germany, Greece, Ireland, Italy, Luxembourg, Netherlands, Portugal, Spain, Sweden, UK

^{†10} countries: Cyprus, Czech Republic, Estonia, Hungary, Latvia, Lithuania, Malta, Poland, Slovakia, Slovenia

[‡]25 countries: listed above for EU15 and EU10

ALL DEVELOPED COUNTRIES: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)		Female (by age)				
Year	0–34	35–69	70+	All	0-34	35–69	70+	All	
1950						•••	•••	•••	
1955	-/878	356/1813 (20%)	90/1638 (5%)	446/4329	-/652	20/1349 (2%)	6.9/2010 (0.3%)	27/4011	
1960	-/736	448/1834 (24%)	137/1799 (8%)	585/4369	-/517	28/1291 (2%)	9.6/2179 (0.4%)	38/3987	
1965	-/663	568/2072 (27%)	217/2007 (11%)	785/4742	-/426	50/1376 (4%)	20/2480 (0.8%)	70/4282	
1970	-/637	662/2268 (29%)	317/2202 (14%)	979/5107	-/372	75/1441 (5%)	42/2804 (1%)	117/4617	
1975	-/605	704/2308 (31%)	420/2432 (17%)	1124/5344	-/340	97/1425 (7%)	73/3132 (2%)	170/4897	
1980	-/597	759/2346 (32%)	492/2682 (18%)	1251/5624	-/312	123/1399 (9%)	112/3501 (3%)	234/5211	
1985	-/555	801/2324 (34%)	552/2918 (19%)	1353/5797	-/299	143/1356 (11%)	170/3941 (4%)	313/5595	
1990	-/521	863/2482 (35%)	546/2862 (19%)	1409/5865	-/258	160/1393 (11%)	235/4023 (6%)	395/5674	
1995	-/518	947/2926 (32%)	594/3102 (19%)	1541/6546	-/238	164/1515 (11%)	302/4417 (7%)	466/6171	
2000	-/425	806/2705 (30%)	634/3322 (19%)	1440/6452	-/190	155/1404 (11%)	349/4619 (8%)	504/6213	

50-year total* (M=millions), mid-1950 to mid-2000: 63/ 517M

-/32M	34/114M	19/121M	52/268M -/20M	4.7/70M	5.7/160M	10/250M
	(30%)	(15%)		(7%)	(4%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: ALL DEVELOPED COUNTRIES



EU15 (European Union - 15 countries): 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


-		Male (l	oy age)		Female (by age)				
Year	0–34	35–69	70+	All	0-34	35–69	70+	All	
1950									
1955	-/235	153/659 (23%)	44/757 (6%)	197/1651	-/173	7.2/495 (1%)	2.6/894 (0.3%)	9.8/1562	
1960	-/205	196/706 (28%)	70/820 (9%)	265/1731	-/141	11/489 (2%)	6.0/992 (0.6%)	17/1622	
1965	-/183	235/780 (30%)	105/872 (12%)	340/1836	-/120	17/499 (3%)	11/1092 (1%)	28/1710	
1970	-/162	249/798 (31%)	149/932 (16%)	398/1892	-/99	20/494 (4%)	18/1198 (2%)	39/1792	
1975	-/136	247/773 (32%)	201/1039 (19%)	448/1947	-/78	23/462 (5%)	27/1313 (2%)	50/1853	
1980	-/117	234/707 (33%)	219/1086 (20%)	453/1910	-/63	27/407 (7%)	35/1365 (3%)	61/1835	
1985	-/97	225/658 (34%)	241/1143 (21%)	465/1898	-/49	30/364 (8%)	52/1461 (4%)	81/1874	
1990	-/97	221/668 (33%)	220/1094 (20%)	441/1859	-/45	33/362 (9%)	62/1462 (4%)	95/1869	
1995	-/85	203/641 (32%)	228/1133 (20%)	431/1859	-/38	33/332 (10%)	81/1511 (5%)	115/1881	
2000	-/66	171/584 (29%)	222/1158 (19%)	394/1808	-/31	34/304 (11%)	89/1531 (6%)	123/1866	

50-year total* (M=millions), mid-1950 to mid-2000: 22/ 181M

-/7.4M	11/35M	8.2/49M	19/92M	-/4.6M	1.1/22M	1.7/63M	2.8/89M
	(30%)	(17%)			(5%)	(3%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

Year



Year

EU10 (European Union - 10 countries): 1960-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)			Female (by age)				
Year	0-34	35–69	70+	AII	0-34	35–69	70+	All		
1950							•••			
1955										
1960										
1965	-/44	38/134 (28%)	14/114 (13%)	52/292	-/28	2.4/94 (3%)	2.8/156 (2%)	5.2/278		
1970	-/41	48/157 (30%)	23/139 (17%)	71/337	-/24	3.1/99 (3%)	3.5/189 (2%)	6.6/311		
1975	-/41	52/161 (32%)	32/163 (19%)	84/365	-/23	3.6/97 (4%)	5.8/215 (3%)	9.4/335		
1980	-/40	66/179 (37%)	39/199 (20%)	106/418	-/20	6.8/101 (7%)	7.3/255 (3%)	14/377		
1985	-/36	76/188 (40%)	40/209 (19%)	116/433	-/18	7.9/99 (8%)	8.6/280 (3%)	16/397		
1990	-/32	91/220 (41%)	34/190 (18%)	124/442	-/14	11/111 (10%)	10/268 (4%)	21/393		
1995	-/26	87/222 (39%)	37/190 (20%)	125/437	-/11	12/107 (11%)	11/275 (4%)	23/393		
2000	-/20	73/196 (37%)	40/191 (21%)	113/407	-/8.0	12/93 (13%)	13/271 (5%)	25/371		


40-year total* (M=millions), mid-1960 to mid-2000: 4.3/ 29M


-/1.5M	2.5/7.0M	1.2/6.8M	3.8/15M	-/0.8M	0.3/4.0M	0.3/9.3M	0.5/14M
	(36%)	(18%)			(6%)	(3%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1965, 1975, 1985 & 1995









EU25 (European Union - 25 countries): 1960-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	oy age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950									
1955									
1960									
1965	-/227	273/914 (30%)	120/987 (12%)	393/2128	-/147	19/593 (3%)	14/1248 (1%)	34/1988	
1970	-/203	296/955 (31%)	172/1071 (16%)	469/2229	-/123	23/593 (4%)	22/1387 (2%)	45/2103	
1975	-/177	299/934 (32%)	232/1202 (19%)	532/2313	-/101	27/559 (5%)	32/1527 (2%)	59/2187	
1980	-/156	300/887 (34%)	258/1285 (20%)	559/2328	-/83	33/508 (7%)	42/1621 (3%)	75/2212	
1985	-/133	301/846 (36%)	281/1353 (21%)	581/2331	-/67	38/463 (8%)	60/1741 (3%)	98/2271	
1990	-/129	312/888 (35%)	253/1283 (20%)	565/2301	-/59	44/473 (9%)	73/1729 (4%)	116/2262	
1995	-/111	290/863 (34%)	266/1323 (20%)	556/2296	-/49	45/439 (10%)	93/1786 (5%)	138/2274	
2000	-/85	244/779 (31%)	263/1350 (19%)	507/2214	-/39	46/397 (11%)	102/1802 (6%)	148/2238	

40-year total* (M=millions), mid-1960 to mid-2000: 24/ 178M

-/6.5M	12/36M	9.0/49M	21/91M	-/3.6M	1.3/21M	2.0/63M	3.3/87M
	(33%)	(18%)			(6%)	(3%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1965, 1975, 1985 & 1995

1965-2000: EU25 (European Union - 25 countries)



AUSTRALIA: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	oy age)			Female (by age)			
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950	-/6.1	2.5/20 (13%)	0.3/18 (2%)	2.8/44	-/4.1	0.0/13	0.0/18	0.0/34	
1955	-/6.1	3.4/21 (17%)	1.3/20 (7%)	4.7/46	-/3.7	0.0/12	0.0/20	0.0/36	
1960	-/6.0	4.7/22 (22%)	1.9/22 (9%)	6.6/50	-/3.8	0.0/13	0.0/22	0.0/39	
1965	-/5.9	6.5/25 (26%)	3.2/25 (13%)	9.7/56	-/3.6	0.1/14 (0.8%)	0.1/27 (0.3%)	0.2/44	
1970	-/6.9	8.4/28 (30%)	4.9/28 (18%)	13/63	-/3.9	1.0/15 (6%)	0.3/31 (0.8%)	1.2/50	
1975	-/6.3	8.6/28 (31%)	5.6/27 (21%)	14/61	-/3.2	1.0/15 (7%)	0.6/30 (2%)	1.6/48	
1980	-/5.6	8.3/26 (32%)	6.6/29 (23%)	15/61	-/2.8	1.5/13 (11%)	1.2/32 (4%)	2.7/48	
1985	-/5.6	7.7/24 (32%)	7.6/34 (22%)	15/64	-/2.8	1.8/13 (14%)	2.0/38 (5%)	3.9/54	
1990	-/5.2	6.7/23 (29%)	7.3/35 (21%)	14/64	-/2.5	1.9/13 (15%)	3.1/40 (8%)	5.0/55	
1995	-/4.6	5.5/22 (26%)	8.1/40 (20%)	14/66	-/2.2	1.8/12 (15%)	4.6/45 (10%)	6.4/59	
2000	-/4.2	4.3/20 (22%)	8.2/43 (19%)	12/67	-/2.0	1.7/11 (15%)	5.0/48 (10%)	6.7/62	


50-year total* (thousands), mid-1950 to mid-2000: 697 / 5337


-/285	319/1190	256/1450	575/2924	-/156	48/663	74/1594	121/2413
	(27%)	(18%)			(7%)	(5%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









AUSTRIA: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All		
1950			•••	•••		•••	•••			
1955	-/5.5	5.6/18 (31%)	2.1/20 (11%)	7.7/44	-/3.6	0.2/14 (2%)	0.2/24 (0.7%)	0.4/41		
1960	-/5.3	6.3/20 (32%)	2.8/21 (13%)	9.0/46	-/3.2	0.3/14 (2%)	0.4/27 (2%)	0.7/43		
1965	-/4.3	6.6/20 (32%)	3.9/23 (17%)	11/47	-/2.7	0.5/14 (3%)	0.4/30 (1%)	0.8/47		
1970	-/4.1	5.9/20 (29%)	4.6/25 (18%)	10/49	-/2.3	0.4/14 (3%)	0.7/34 (2%)	1.1/50		
1975	-/3.5	4.7/17 (27%)	5.2/26 (20%)	9.9/47	-/1.8	0.6/12 (5%)	0.8/36 (2%)	1.4/49		
1980	-/3.0	4.2/15 (27%)	5.1/26 (20%)	9.3/44	-/1.4	0.5/9.9 (5%)	1.4/37 (4%)	1.9/48		
1985	-/2.6	3.9/14 (28%)	4.7/25 (18%)	8.6/42	-/1.1	0.5/8.4 (5%)	1.5/38 (4%)	2.0/48		
1990	-/2.1	4.0/14 (28%)	3.6/22 (16%)	7.6/38	-/0.9	0.6/8.3 (7%)	1.5/35 (4%)	2.0/45		
1995	-/1.9	3.7/14 (27%)	3.5/22 (16%)	7.1/38	-/0.8	0.6/7.1 (8%)	1.5/36 (4%)	2.0/44		
2000	-/1.4	3.1/12 (26%)	3.2/22 (15%)	6.3/35	-/0.6	0.7/6.3 (12%)	1.8/35 (5%)	2.6/42		


50-year total* (thousands), mid-1950 to mid-2000: 506 / 4461


-/178	245/837	195/1158	440/2173	-/100	23/551	44/1637	67/2288
	(29%)	(17%)			(4%)	(3%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









BELARUS: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All		
1950						•••				
1955						•••	•••			
1960										
1965										
1970										
1975										
1980										
1985	-/5.6	8.5/24 (36%)	2.4/22 (11%)	11/51	-/2.6	0.0/14	0.0/38	0.0/55		
1990	-/5.4	11/29 (37%)	2.8/20 (14%)	13/53	-/2.1	0.2/16	0.1/38	0.3/56		
1995	-/5.8	15/40 (36%)	3.9/22 (17%)	19/69	-/2.1	0.0/20	0.0/43	0.0/65		
2000	-/5.3	13/39 (33%)	4.7/26 (18%)	18/70	-/1.8	0.0/18	0.0/45	0.0/65		

20-year total* (thousands), mid-1980 to mid-2000: 294 / 2395

-/114	231/643	63/440	294/1197	-/46	0.0/337	0.0/814	0.0/1198
	(36%)	(14%)			(0%)	(0%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: BELARUS



BELGIUM: 1950-2000[‡]

[‡]2000 mortality involves 1997 rates applied to 2000 population

Numbers of deaths attributed to smoking / total deaths (thousands)

		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950			•••	•••				•••		
1955	-/6.2	6.4/24 (27%)	1.5/28 (6%)	7.9/58	-/4.2	0.0/16	0.0/31	0.0/51		
1960	-/5.3	8.5/26 (33%)	2.5/30 (8%)	11/61	-/3.4	0.0/16	0.0/34	0.0/53		
1965	-/4.6	10/27 (37%)	4.3/30 (14%)	14/62	-/2.9	0.1/15 (0.5%)	0.1/35 (0.4%)	0.2/53		
1970	-/4.2	10/26 (39%)	6.0/32 (19%)	16/63	-/2.5	0.1/15 (0.9%)	0.1/38 (0.1%)	0.2/56		
1975	-/3.6	9.8/24 (41%)	8.4/35 (24%)	18/63	-/1.9	0.4/14 (3%)	0.6/41 (1%)	1.0/57		
980	-/3.5	9.2/22 (42%)	9.3/34 (27%)	18/60	-/1.9	0.5/12 (4%)	0.6/41 (1%)	1.1/54		
985	-/2.8	8.7/20 (44%)	10/35 (29%)	19/57	-/1.4	0.5/10 (5%)	0.8/43 (2%)	1.4/54		
1990	-/2.7	7.8/19 (41%)	8.9/31 (28%)	17/53	-/1.3	0.8/10 (8%)	0.8/40 (2%)	1.6/51		
1995	-/2.3	7.4/18 (41%)	9.6/33 (29%)	17/53	-/1.1	1.0/9.6 (10%)	1.5/41 (4%)	2.4/52		
2000	-/2.0	6.5/17 (39%)	9.4/34 (28%)	16/52	-/1.0	0.9/8.8 (10%)	1.8/43 (4%)	2.7/53		


50-year total* (thousands), mid-1950 to mid-2000: 813 / 5597


-/194	423/1122	340/1610	763/2926	-/116	20/646	30/1909	50/2671
	(38%)	(21%)			(3%)	(2%)	


 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995


1955-2000[‡]: BELGIUM

[‡]2000 mortality involves 1997 rates applied to 2000 population









BULGARIA: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950						•••	•••			
955										
960										
965										
970										
975										
980										
985	-/3.4	7.5/24 (31%)	2.8/32 (9%)	10/59	-/2.0	0.7/13 (6%)	0.9/33 (3%)	1.6/48		
990	-/3.3	8.4/27 (31%)	2.2/30 (8%)	11/60	-/1.8	0.5/14 (4%)	0.5/33 (1%)	1.0/49		
995	-/2.8	8.8/28 (31%)	2.7/32 (8%)	12/63	-/1.5	0.8/14 (6%)	0.9/36 (2%)	1.7/51		
2000	-/2.3	7.4/25 (29%)	2.6/34 (8%)	10/62	-/1.3	0.7/13 (5%)	0.5/39 (1%)	1.2/54		

20-year total* (thousands), mid-1980 to mid-2000: 251 / 2222

-/62	163/524	55/641	218/1227	-/35	15/264	18/696	33/994
	(31%)	(9%)			(6%)	(3%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: BULGARIA



CANADA: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)

		Male (I	oy age)			Female (by age)				
Year	0-34	35–69	70+	AII	0-34	35–69	70+	All		
1950	-/15	3.5/28 (13%)	0.5/27 (2%)	4.0/70	-/11	0.0/18	0.0/25	0.0/53		
1955	-/14	4.9/29 (17%)	1.4/32 (4%)	6.3/75	-/9.4	0.1/17	0.0/27	0.1/53		
1960	_/14	6.3/31 (20%)	2.6/37 (7%)	8.9/82	-/8.8	0.0/18	0.0/31	0.0/58		
1965	-/12	8.4/34 (25%)	3.9/41 (10%)	12/87	-/7.5	0.5/19 (3%)	0.0/36	0.5/62		
1970	_/11	11/38 (28%)	5.8/42 (14%)	17/91	-/6.4	1.0/20 (5%)	0.3/38 (0.7%)	1.3/65		
1975	_/11	12/40 (30%)	8.0/45 (18%)	20/97	-/5.7	2.2/22 (10%)	0.7/43 (2%)	2.9/70		
1980	-/10	13/40 (33%)	9.9/48 (21%)	23/98	-/4.9	3.1/22 (14%)	2.3/47 (5%)	5.4/74		
1985	-/8.2	13/38 (35%)	12/54 (23%)	25/100	-/4.0	4.5/22 (21%)	4.5/55 (8%)	9.0/81		
1990	-/7.7	13/38 (35%)	14/58 (24%)	27/104	-/3.7	5.0/22 (23%)	6.9/62 (11%)	12/88		
1995	-/6.9	12/38 (31%)	16/66 (24%)	27/111	-/3.3	5.6/23 (25%)	10/74 (14%)	16/99		
2000	-/5.1	10/36 (27%)	16/70 (22%)	26/112	-/2.7	5.9/22 (26%)	13/81 (16%)	19/106		


50-year total* (thousands), mid-1950 to mid-2000: 1200 / 8363

-/524	505/1797	412/2386	917/4707	-/299	128/1013	154/2344	283/3656
	(28%)	(17%)			(13%)	(7%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1950-2000: CANADA

Year



Year

See note on page 127

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950		•••								
1955				•••						
1960										
1965										
1970										
1975										
1980										
1985	-/101	34/105 (32%)	8.6/83 (10%)	43/290	-/77	5.0/73 (7%)	3.1/125 (3%)	8.1/275		
1990	-/90	39/126 (31%)	8.9/75 (12%)	48/291	-/62	5.9/80 (7%)	3.9/125 (3%)	9.8/268		
1995	-/87	45/171 (27%)	7.8/84 (9%)	53/341	-/58	4.7/101 (5%)	3.1/147 (2%)	7.7/307		
2000	-/62	35/155 (23%)	9.6/95 (10%)	45/312	-/38	3.9/92 (4%)	3.7/152 (2%)	7.6/282		

20-year total* (thousands), mid-1980 to mid-2000: 1117 / 12134

-/1886	795/2761	164/1667	958/6313	-/1355	96/1741	62/2724	159/5820
	(29%)	(10%)			(6%)	(2%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: CENTRAL ASIA (8 countries)



CZECH REPUBLIC: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	AII	0-34	35–69	70+	All		
1950			•••			•••	•••	•••		
955										
960			•••							
965		•••	•••			•••	•••	•••		
1970			•••				•••	•••		
975			•••			•••	•••	•••		
980			•••				•••	•••		
985	-/3.3	12/27 (43%)	7.9/36 (22%)	20/67	-/1.6	1.0/14 (7%)	1.4/49 (3%)	2.5/65		
990	-/3.0	13/31 (42%)	6.1/32 (19%)	19/66	-/1.5	1.4/16 (9%)	1.5/45 (3%)	2.9/63		
995	-/2.5	10/26 (39%)	6.1/31 (20%)	16/59	-/1.2	1.6/14 (12%)	1.5/44 (3%)	3.1/59		
2000	-/2.0	8.5/23 (37%)	5.6/30 (19%)	14/55	-/0.8	1.4/12 (12%)	2.3/42 (5%)	3.6/54		

20-year total* (thousands), mid-1980 to mid-2000: 414 / 2496

-/58	218/528	140/669	358/1255	-/28	26/278	29/934	55/1240
	(41%)	(21%)			(9%)	(3%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: CZECH REPUBLIC



DENMARK: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All		
1950				•••			•••			
1955	-/2.2	1.4/7.8 (18%)	0.4/10 (4%)	1.8/20	-/1.5	0.0/6.3	0.0/11	0.0/18		
1960	-/2.0	2.0/8.6 (23%)	0.8/12 (6%)	2.7/23	-/1.2	0.1/6.4 (2%)	0.3/13 (2%)	0.4/21		
1965	-/2.0	2.9/10 (28%)	1.4/14 (10%)	4.3/26	-/1.3	0.3/6.7 (4%)	0.3/14 (2%)	0.5/22		
1970	-/1.7	3.0/10 (30%)	1.9/14 (13%)	4.9/26	-/1.0	0.5/6.8 (7%)	0.5/14 (3%)	1.0/22		
1975	-/1.5	3.3/10 (31%)	2.7/16 (17%)	5.9/28	-/0.8	0.7/6.5 (10%)	0.6/16 (4%)	1.3/23		
1980	-/1.4	3.6/11 (32%)	3.8/18 (21%)	7.3/30	-/0.7	1.0/6.7 (15%)	1.1/18 (6%)	2.1/26		
1985	-/1.3	3.6/11 (34%)	4.2/18 (23%)	7.9/30	-/0.6	1.5/6.6 (22%)	1.7/21 (8%)	3.2/28		
1990	-/1.2	3.3/10 (32%)	4.3/20 (22%)	7.6/31	-/0.6	1.8/6.8 (27%)	2.6/22 (11%)	4.4/30		
1995	-/1.1	2.9/9.8 (30%)	4.6/20 (23%)	7.5/31	-/0.5	2.0/6.8 (30%)	3.4/24 (14%)	5.4/32		
2000	-/0.8	2.5/8.8 (28%)	4.1/18 (23%)	6.6/28	-/0.4	1.8/6.1 (29%)	4.0/23 (18%)	5.8/29		

50-year total* (thousands), mid-1950 to mid-2000: 379 / 2582

- /81	141/487		274/1352	-/47	44/329	60/854	105/1230
	(29%)	(17%)			(13%)	(7%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: DENMARK



ESTONIA: 1980-2000

Numbers of deaths attributed to smoking / total deaths


		Male (b	y age)		Female (by age)				
Year	0–34	35–69	70+	All	0–34	35–69	70+	All	
1950			•••				•••		
1955									
1960									
1965									
1970			•••				•••	•••	
1975							•••	•••	
1980							•••	•••	
1985	-/901	1550/4286 (36%)	617/3911 (16%)	2167/9098	-/373	93/2403 (4%)	132/7469 (2%)	225/10245	
1990	-/930	1848/4891 (38%)	529/3603 (15%)	2377/9424	-/355	172/2666 (6%)	113/7085 (2%)	285/10106	
1995	-/918	2188/6383 (34%)	611/3532 (17%)	2799/10833	-/291	157/2854 (6%)	103/6850 (2%)	260/9995	
2000	-/546	1557/4963 (31%)	763/3756 (20%)	2320/9265	-/202	77/2315 (3%)	354/6621 (5%)	431/9138	

20-year total* (thousands), mid-1980 to mid-2000: 55 / 402

-/18	37/107	12/74	50/199	-/6.6	2.5/53	2.3/143	4.8/202
	(35%)	(16%)			(5%)	(2%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: ESTONIA



FINLAND: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All		
1950				•••						
1955	-/3.5	3.5/10 (34%)	0.8/6.7 (12%)	4.3/20	-/2.2	0.1/6.6	0.0/10	0.1/19		
1960	-/2.7	4.1/11 (38%)	1.1/7.2 (16%)	5.2/21	-/1.5	0.0/6.5	0.0/11	0.0/19		
1965	-/2.4	4.8/12 (39%)	1.7/8.5 (20%)	6.5/23	-/1.3	0.0/7.0	0.0/13	0.0/21		
1970	-/2.1	4.9/13 (38%)	2.1/9.0 (23%)	6.9/24	-/1.0	0.0/6.7	0.0/13	0.0/20		
1975	-/2.0	4.2/12 (35%)	2.2/9.7 (23%)	6.4/24	-/0.8	0.0/6.1	0.1/13 (0.5%)	0.1/20		
1980	-/1.7	3.8/11 (35%)	3.0/11 (27%)	6.8/24	-/0.7	0.2/5.2 (4%)	0.4/15 (3%)	0.6/21		
1985	-/1.4	3.4/11 (32%)	3.1/13 (23%)	6.4/25	-/0.6	0.3/4.9 (6%)	0.4/18 (2%)	0.7/23		
1990	-/1.4	2.6/11 (25%)	2.7/13 (21%)	5.3/25	-/0.6	0.2/4.9 (5%)	0.5/20 (3%)	0.8/25		
1995	-/1.1	2.1/9.9 (22%)	2.6/13 (20%)	4.8/24	-/0.4	0.2/4.4 (5%)	0.6/20 (3%)	0.8/25		
2000	-/0.9	1.6/9.1 (17%)	2.6/14 (18%)	4.1/24	-/0.4	0.2/4.2 (5%)	0.8/21 (4%)	1.0/25		


50-year total* (thousands), mid-1950 to mid-2000: 301 / 2256


-/104	180/553	104/512	284/1169	-/53	6.7/289	10/745	17/1087
	(33%)	(20%)			(2%)	(1%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









FRANCE: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950	-/41	11/111 (10%)	1.0/117 (0.9%)	12/270	-/30	0.0/84	0.0/146	0.0/261	
1955	-/31	17/112 (15%)	3.8/123 (3%)	21/267	-/21	0.0/77	0.0/158	0.0/256	
1960	-/24	22/114 (20%)	6.1/125 (5%)	29/264	-/16	0.0/70	0.0/168	0.0/254	
1965	-/23	28/127 (22%)	9.7/128 (8%)	38/279	-/14	0.0/68	0.0/179	0.0/262	
1970	-/23	29/122 (24%)	14/133 (10%)	43/278	-/13	0.0/63	0.0/186	0.0/262	
1975	-/21	33/119 (28%)	21/151 (14%)	53/291	-/11	0.0/58	0.0/200	0.0/269	
1980	-/21	33/106 (31%)	23/158 (14%)	55/285	-/10	0.0/48	0.0/203	0.0/262	
1985	-/18	33/102 (32%)	26/167 (16%)	59/287	-/8.9	0.5/44 (1%)	0.9/213 (0.4%)	1.4/266	
1990	-/18	34/103 (33%)	24/152 (16%)	58/273	-/7.7	1.2/44 (3%)	1.3/201 (0.6%)	2.5/254	
1995	-/15	33/100 (34%)	25/161 (16%)	59/275	-/6.8	2.0/43 (5%)	2.5/206 (1%)	4.6/257	
2000	-/11	30/91 (32%)	25/169 (15%)	54/272	-/5.3	2.5/41 (6%)	3.4/213 (2%)	5.9/259	


50-year total* (M=millions), mid-1950 to mid-2000: 2.4/ 27M


-/1.1M	1.4/5.6M	0.9/7.3M	2.3/14M	-/0.6M	0.02/2.9M	0.03/9.6M	0.06/13M
	(26%)	(12%)			(0.8%)	(0.4%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









GERMANY: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950			•••	•••			•••		
1955	-/50	34/153 (22%)	9.0/181 (5%)	43/384	-/36	0.2/129 (0.1%)	0.0/207	0.2/371	
1960	-/48	47/173 (27%)	17/208 (8%)	64/430	-/31	1.5/135 (1%)	0.3/245 (0.1%)	1.8/411	
1965	_/41	57/193 (29%)	28/225 (12%)	84/459	-/25	3.1/142 (2%)	3.7/282 (1%)	6.9/449	
1970	-/38	60/200 (30%)	42/245 (17%)	102/483	-/21	2.9/143 (2%)	3.2/329 (1%)	6.1/493	
1975	-/30	54/180 (30%)	56/271 (21%)	109/481	-/16	2.8/129 (2%)	4.4/364 (1%)	7.1/508	
1980	-/25	48/156 (31%)	55/274 (20%)	104/456	-/13	3.1/107 (3%)	6.1/376 (2%)	9.2/496	
1985	-/20	46/141 (33%)	54/272 (20%)	100/434	-/10	4.1/89 (5%)	7.2/396 (2%)	11/496	
1990	-/20	52/162 (32%)	43/243 (18%)	95/425	-/9.3	6.2/94 (7%)	10/393 (3%)	17/496	
1995	-/16	51/165 (31%)	43/230 (19%)	94/411	-/7.5	7.1/84 (8%)	16/383 (4%)	23/474	
2000	-/12	43/149 (29%)	41/227 (18%)	84/389	-/5.8	8.1/75 (11%)	16/369 (4%)	24/450	


50-year total* (M=millions), mid-1950 to mid-2000: 4.8/ 45M


-/1.6M	2.4/8.3M	1.9/12M	4.3/22M	-/0.9M	0.2/5.7M	0.3/16M	0.5/23M
	(29%)	(16%)			(3%)	(2%)	


 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: GERMANY









GREECE: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950				•••						
1955	-/6.7	2.0/9.6 (21%)	0.7/11 (6%)	2.7/28	-/5.8	0.5/7.8 (6%)	0.0/13	0.5/27		
1960	-/6.0	2.4/11 (23%)	1.0/14 (7%)	3.4/31	-/4.8	0.3/7.7 (4%)	0.0/17	0.4/30		
1965	-/5.1	3.1/13 (24%)	1.6/17 (10%)	4.7/34	-/3.8	0.5/9.1 (5%)	0.1/20 (0.3%)	0.5/33		
1970	-/4.3	3.5/14 (24%)	2.6/20 (13%)	6.1/38	-/3.0	0.5/9.5 (5%)	0.2/23 (0.9%)	0.7/36		
1975	-/3.8	4.2/15 (29%)	3.9/23 (17%)	8.1/41	-/2.5	0.6/9.4 (6%)	0.7/27 (3%)	1.2/39		
1980	-/3.3	4.5/15 (30%)	4.4/27 (16%)	8.8/45	-/2.0	0.3/9.2 (3%)	0.5/31 (2%)	0.8/42		
1985	-/2.9	4.8/15 (32%)	5.5/31 (18%)	10/48	-/1.5	0.5/8.8 (5%)	0.7/34 (2%)	1.2/44		
1990	-/2.3	5.1/16 (33%)	5.4/32 (17%)	11/49	-/1.3	0.4/8.2 (5%)	1.0/35 (3%)	1.4/45		
1995	-/2.4	5.6/17 (33%)	5.8/33 (17%)	11/53	-/1.0	0.4/8.6 (5%)	1.0/38 (3%)	1.4/47		
2000	-/2.2	5.5/17 (33%)	6.6/37 (18%)	12/55	-/0.9	0.4/7.9 (5%)	1.3/41 (3%)	1.7/50		


50-year total* (thousands), mid-1950 to mid-2000: 420 / 3952


-/209	197/690	176/1150	372/2048	-/146	24/437	24/1321	48/1903
	(28%)	(15%)			(5%)	(2%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









HUNGARY: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950				•••		•••			
1955	_/11	3.9/20 (20%)	0.7/19 (4%)	4.6/50	-/8.2	0.6/17 (3%)	0.1/23 (0.6%)	0.7/48	
1960	-/7.3	4.9/22 (23%)	1.6/23 (7%)	6.5/52	-/5.1	0.7/17 (4%)	0.5/27 (2%)	1.2/50	
1965	-/5.8	6.3/24 (26%)	2.7/26 (11%)	9.1/55	-/3.8	0.9/17 (5%)	0.9/31 (3%)	1.8/53	
1970	-/6.2	7.1/27 (26%)	4.1/29 (14%)	11/63	-/3.7	1.1/18 (6%)	1.1/36 (3%)	2.1/58	
1975	-/6.7	8.0/29 (28%)	6.3/33 (19%)	14/68	-/4.1	1.0/19 (5%)	1.9/40 (5%)	2.8/63	
1980	-/5.2	11/33 (34%)	8.3/38 (22%)	20/77	-/2.9	1.9/20 (10%)	2.4/46 (5%)	4.4/69	
1985	-/4.8	13/34 (38%)	8.1/39 (21%)	21/78	-/2.5	2.1/19 (11%)	2.8/48 (6%)	4.9/70	
1990	-/4.2	16/39 (41%)	6.5/34 (19%)	23/77	-/2.1	3.0/21 (14%)	3.1/46 (7%)	6.0/69	
1995	-/3.2	16/40 (41%)	7.5/34 (22%)	24/77	-/1.5	3.5/20 (18%)	3.5/47 (7%)	7.0/68	
2000	-/2.5	14/35 (41%)	7.1/33 (21%)	21/70	-/1.2	3.7/17 (21%)	3.6/46 (8%)	7.3/65	

50-year total* (thousands), mid-1950 to mid-2000: 899 / 6301

-/318	474/1465	253/1505	726/3289	-/201	81/919	92/1892	172/3012
	(32%)	(17%)			(9%)	(5%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: HUNGARY



Year

Year

IRELAND: 1950-2000


Numbers of deaths attributed to smoking / total deaths


		Male (k	oy age)		Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All	
1950	-/3138	800/6938 (12%)	15/9950 (0.2%)	815/20026	-/2716	0/5527	0/9472	0/17715	
1955	-/2209	1337/6623 (20%)	245/10941 (2%)	1582/19773	-/1702	313/4869 (6%)	0/10417	313/16988	
1960	-/1718	1633/6026 (27%)	358/9740 (4%)	1991/17484	-/124 4	312/4337 (7%)	7/9595	319/15176	
1965	-/1533	1938/6510 (30%)	776/9922 (8%)	2714/17965	-/1087	361/4106 (9%)	162/9864 (2%)	523/15057	
1970	-/1365	2437/6980 (35%)	1222/10102 (12%)	3659/18447	-/970	585/4317 (14%)	429/9952 (4%)	1014/15239	
1975	-/1450	2274/6779 (34%)	1720/9848 (17%)	3994/18077	-/8 7 5	700/4037 (17%)	628/10184 (6%)	1328/15096	
1980	-/1230	2174/6794 (32%)	2027/10206 (20%)	4201/18230	-/ 7 35	750/4004 (19%)	752/10503 (7%)	1502/15242	
1985	-/1048	2180/6228 (35%)	2507/10925 (23%)	4687/18201	-/562	732/3589 (20%)	1469/10861 (14%)	2201/15012	
1990	-/883	1670/5475 (31%)	2476/10470 (24%)	4146/16828	-/498	652/3245 (20%)	1607/10799 (15%)	2259/14542	
1995	-/878	1448/5144 (28%)	2506/11053 (23%)	3954/17075	-/447	610/3076 (20%)	1627/11661 (14%)	2237/15184	
2000	-/979	1182/4902 (24%)	2125/10311 (21%)	3307/16192	-/432	518/2953 (18%)	1904/11814 (16%)	2422/15199	


50-year total* (thousands), mid-1950 to mid-2000: 235 / 1684


-/71	92/313	78/527	169/911	-/47	27/197	39/530	66/773
	(29%)	(15%)			(14%)	(7%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









ITALY: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)			Female (by age)				
Year	0–34	35–69	70+	All	0–34	35–69	70+	All		
1950			•••	•••		•••		•••		
1955	-/45	14/88 (15%)	1.1/102 (1%)	15/234	-/34	0.0/67	0.0/112	0.0/212		
1960	-/40	22/99 (22%)	3.1/116 (3%)	25/255	-/29	0.6/68 (0.8%)	0.1/129 (0.1%)	0.7/226		
1965	-/35	30/113 (26%)	7.5/128 (6%)	37/275	-/24	1.1/70 (2%)	0.3/148 (0.2%)	1.4/243		
1970	-/29	33/115 (29%)	14/133 (10%)	47/277	-/19	1.2/68 (2%)	0.7/157 (0.4%)	1.9/244		
1975	-/22	39/121 (32%)	23/152 (15%)	62/295	-/13	1.7/66 (3%)	3.1/180 (2%)	4.8/259		
1980	-/17	41/115 (36%)	30/162 (18%)	71/294	-/9.5	2.3/60 (4%)	3.5/190 (2%)	5.7/260		
1985	-/13	39/103 (38%)	36/171 (21%)	75/287	-/6.8	2.5/53 (5%)	6.6/201 (3%)	9.1/261		
1990	-/14	38/102 (37%)	35/166 (21%)	72/282	-/6.0	2.7/53 (5%)	7.5/203 (4%)	10/262		
1995	-/13	32/94 (34%)	39/180 (22%)	71/286	-/5.6	2.8/48 (6%)	8.6/217 (4%)	11/270		
2000	-/9.1	25/81 (31%)	42/191 (22%)	66/281	-/4.1	2.7/43 (6%)	11/232 (5%)	13/279		


50-year total* (M=millions), mid-1950 to mid-2000: 2.9/ 26M


-/1.3M	1.5/5.2M	1.1/7.3M	2.6/14M	-/0.8M	0.1/3.0M	0.2/8.6M	0.3/12M
	(30%)	(15%)			(3%)	(2%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995









JAPAN: 1950-2000


Numbers of deaths attributed to smoking / total deaths (thousands)

		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	AII	
1950	-/192	0.0/177 (0%)	0.0/98 (0%)	0.0/467	-/178	0.0/140	0.0/120	0.0/438	
1955	-/102	2.7/156 (2%)	0.7/107 (0.7%)	3.5/365	-/84	0.0/115	0.0/129	0.0/328	
1960	-/77	9.2/165 (6%)	3.4/135 (2%)	13/378	-/57	0.0/114	0.0/159	0.0/329	
1965	-/56	13/169 (8%)	6.6/153 (4%)	19/379	-/37	1.9/111 (2%)	1.2/174 (0.7%)	3.2/322	
1970	-/49	17/170 (10%)	12/169 (7%)	29/388	-/30	2.7/111 (2%)	2.6/185 (1%)	5.3/325	
1975	-/38	18/156 (11%)	18/184 (10%)	36/378	-/24	3.1/101 (3%)	5.2/200 (3%)	8.3/324	
1980	-/28	20/154 (13%)	26/209 (13%)	47/391	-/16	3.7/95 (4%)	9.7/221 (4%)	13/332	
1985	-/22	22/157 (14%)	35/229 (15%)	57/408	-/12	3.8/90 (4%)	12/242 (5%)	16/345	
1990	-/19	27/168 (16%)	42/257 (16%)	68/444	-/10	3.6/88 (4%)	16/278 (6%)	19/377	
1995	-/18	31/188 (16%)	53/296 (18%)	84/501	-/9.7	4.4/91 (5%)	20/320 (6%)	25/421	
2000	-/15	29/184 (16%)	61/327 (19%)	90/526	-/8.1	4.2/85 (5%)	20/343 (6%)	24/436	

50-year total* (M=millions), mid-1950 to mid-2000: 2.5/ 38M

	(10%)	(12%)			(3%)	(4%)	
-/2.4M	0.9/8.3M	1.1/9.7M	2.0/20M	-/1.7M	0.1/5.1M	0.4/11M	0.5/17M

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995



LATVIA: 1980-2000


Numbers of deaths attributed to smoking / total deaths


		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	AII	
1950			•••	•••		•••		•••	
1955									
1960									
1965									
1970									
1975									
1980									
1985	-/1500	2735/7222 (38%)	849/7239 (12%)	3584/15961	-/661	236/4417 (5%)	79/13127 (0.6%)	315/18205	
1990	-/1702	3180/8526 (37%)	997/6723 (15%)	4177/16951	-/656	274/4816 (6%)	242/12389 (2%)	516/17861	
1995	-/1749	3874/12037 (32%)	1095/6557 (17%)	4969/20343	-/584	197/5616 (4%)	216/12388 (2%)	413/18588	
2000	-/1088	2663/8848 (30%)	1066/6219 (17%)	3729/16155	-/373	76/4107 (2%)	348/11570 (3%)	424/16050	


20-year total* (thousands), mid-1980 to mid-2000: 93 / 731


-/32	66/193	19/138	86/363	-/12	4.3/100	3.0/255	7.3/368
	(34%)	(14%)			(4%)	(1%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995









LITHUANIA: 1980-2000

Numbers of deaths attributed to smoking / total deaths


		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	AII	0–34	35–69	70+	AII	
1950		•••	•••	•••			•••		
1955									
1960									
1965									
1970									
1975									
1980									
1985	-/2236	3044/8678 (35%)	1641/9370 (18%)	4685/20284	-/914	204/4772	194/13199	398/18885	
1990	-/1958	3789/10184 (37%)	1441/8463 (17%)	5230/20605	-/773	176/5391	428/12991	604/19155	
1995	-/2200	4549/13545 (34%)	1524/8538 (18%)	6073/24283	-/783	160/6324	287/13916	447/21023	
2000	-/1610	3038/10401 (29%)	1614/8397 (19%)	4652/20408	-/568	0/4794	0/13149	0/18511	

20-year total* (thousands), mid-1980 to mid-2000: 116 / 845

-/44	76/222	32/179	108/446	-/17	3.6/111	4.8/271	8.4/399
	(34%)	(18%)			(3%)	(2%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: LITHUANIA



LUXEMBOURG: 1950-2000

Numbers of deaths attributed to smoking / total deaths

		Male (b	y age)		Female (by age)				
Year	0–34	35–69	70+	AII	0-34	35–69	70+	All	
1950									
1955	-/223	232/856 (27%)	40/833 (5%)	272/1912	-/146	0/591	0/858	0/1595	
1960	-/211	279/893 (31%)	61/921 (7%)	340/2025	-/108	0/571	0/1012	0/1691	
1965	-/168	415/1049 (40%)	145/1030 (14%)	560/2247	-/94	0/593	0/1084	0/1771	
1970	-/181	432/1021 (42%)	175/1089 (16%)	607/2291	-/107	9/627	0/1131	9/1865	
1975	-/156	402/1032 (39%)	250/1144 (22%)	652/2332	-/62	2/576	20/1405	22/2043	
1980	-/105	321/827 (39%)	261/1184 (22%)	582/2116	-/70	22/483	10/1465	32/2018	
1985	-/107	283/766 (37%)	315/1227 (26%)	598/2100	-/61	41/443	0/1476	41/1980	
1990	-/108	249/727 (34%)	274/1078 (25%)	523/1913	-/51	41/443	21/1426	62/1920	
1995	-/90	263/766 (34%)	272/1043 (26%)	535/1899	-/47	13/356	97/1398 (7%)	110/1801	
2000	-/96	209/667 (31%)	236/1094 (22%)	445/1857	-/39	40/354	85/1459 (6%)	125/1852	


50-year total* (thousands), mid-1950 to mid-2000: 28 / 197

-/7.4	16/45	10/53	26/105	-/4.1	0.6/26	1.2/62	1.7/92
	(36%)	(19%)			(2%)	(2%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: LUXEMBOURG

Year



Year

MALTA: 1960-2000

Numbers of deaths attributed to smoking / total deaths


		Male (k	oy age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950									
1955									
1960									
1965	-/173	205/657 (31%)	70/739 (9%)	275/1569	-/134	4/469	0/829	4/1432	
1970	-/156	218/688 (32%)	65/775 (8%)	283/1619	-/89	0/467	0/895	0/1451	
1975	-/101	252/643 (39%)	68/827 (8%)	320/1571	-/90	0/424	0/924	0/1438	
1980	-/123	197/625 (32%)	90/964 (9%)	287/1712	-/67	0/448	0/1074	0/1589	
1985	-/117	176/562 (31%)	136/826 (16%)	312/1505	-/71	0/330	0/1009	0/1410	
1990	-/71	141/484 (29%)	161/836 (19%)	302/1391	-/36	0/325	0/957	0/1318	
1995	-/86	141/442 (32%)	171/839 (20%)	312/1367	-/32	0/289	0/1020	0/1341	
2000	-/49	116/474 (24%)	127/990 (13%)	243/1513	-/32	10/318	41/1110	51/1460	

40-year total* (thousands), mid-1960 to mid-2000: 12 / 116

-/4.8	7.7/23	4.4/32	12/60	-/3.3	0.0/15	0.0/38	0.0/56
	(34%)	(14%)			(0.3%)	(0%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1965, 1975, 1985 & 1995

1965-2000: MALTA



MOLDOVA, Republic of: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950				•••		•••		•••		
1955										
1960										
1965										
1970										
1975										
1980										
1985	-/3.9	3.2/11 (29%)	0.7/8.1 (9%)	3.9/23	-/2.2	0.4/9.0 (5%)	0.1/12	0.5/23		
1990	-/3.0	3.5/11 (31%)	0.7/7.3 (10%)	4.2/21	-/1.5	0.3/8.4 (3%)	0.3/11	0.5/21		
1995	-/2.8	4.3/15 (29%)	0.8/9.4 (9%)	5.1/27	-/1.4	0.4/10 (4%)	0.0/14	0.5/26		
2000	-/1.8	2.6/11 (23%)	0.5/8.2 (7%)	3.1/21	-/0.8	0.2/7.4 (3%)	0.0/12	0.2/20		

20-year total* (thousands), mid-1980 to mid-2000: 99 / 989

-/68	75/258	15/175	90/501	-/36	8.6/192	0.9/261	9.5/489
	(29%)	(9%)			(4%)	(0.3%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: Republic of MOLDOVA



NETHERLANDS: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	by age)		Female (by age)					
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950	-/6.6	3.0/14 (22%)	0.9/18 (5%)	3.9/39	-/4.7	0.0/12	0.0/20	0.0/37		
1955	-/5.4	4.3/16 (27%)	1.5/22 (7%)	5.8/43	-/3.7	0.0/12	0.0/22	0.0/38		
1960	-/5.0	6.0/18 (34%)	2.6/25 (10%)	8.6/47	-/3.3	0.0/12	0.0/25	0.0/40		
1965	-/4.9	8.2/21 (39%)	4.3/29 (15%)	12/54	-/3.1	0.0/13	0.0/28	0.0/44		
1970	-/5.0	9.9/24 (42%)	6.7/32 (21%)	17/61	-/3.0	0.0/13	0.0/32	0.0/49		
1975	-/3.9	10/24 (44%)	9.2/36 (26%)	20/64	-/2.3	0.1/13 (0.7%)	0.0/35	0.1/50		
1980	-/3.6	9.8/23 (43%)	11/37 (31%)	21/63	-/2.0	0.5/12 (4%)	0.2/37 (0.7%)	0.8/51		
1985	-/3.1	9.8/23 (43%)	13/40 (32%)	23/66	-/1.8	0.9/12 (8%)	0.7/43 (2%)	1.6/57		
1990	-/3.0	8.6/22 (38%)	13/41 (32%)	22/67	-/1.8	1.4/12 (11%)	1.3/48 (3%)	2.7/62		
1995	-/2.6	7.9/22 (36%)	13/44 (29%)	21/68	-/1.5	2.2/13 (17%)	2.9/53 (5%)	5.1/67		
2000	-/2.3	6.7/22 (31%)	12/45 (27%)	19/69	-/1.4	2.8/13 (21%)	4.2/57 (7%)	7.0/72		


50-year total* (thousands), mid-1950 to mid-2000: 880 / 5515


-/199	406/1056	406/1698	812/2954	-/123	32/621	36/1817	68/2562
	(38%)	(24%)			(5%)	(2%)	


 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1950-2000: NETHERLANDS









NEW ZEALAND: 1950-2000

Numbers of deaths attributed to smoking / total deaths


		Male (k	oy age)		Female (by age)				
Year	0–34	35–69	70+	All	0–34	35–69	70+	All	
1950	-/1542	569/3968 (14%)	145/4431 (3%)	714/9941	-/1204	0/2941	0/3998	0/8143	
1955	-/1609	932/3998 (23%)	370/5020 (7%)	1302/10627	-/1022	0/2831	0/4745	0/8598	
1960	-/1539	970/4294 (23%)	578/5713 (10%)	1548/11546	-/1042	58/2734 (2%)	17/5570	75/9346	
1965	-/1529	1408/5125 (27%)	911/6005 (15%)	2319/12659	-/915	95/3121 (3%)	0/6281	95/10317	
1970	-/1576	1909/5822 (33%)	1114/6217 (18%)	3023/13615	-/916	321/3366 (10%)	199/6943 (3%)	520/11225	
1975	-/1560	1924/5902 (33%)	1422/6336 (22%)	3346/13798	-/902	454/3370 (13%)	309/7045 (4%)	763/11317	
1980	-/1367	1776/5617 (32%)	1755/7354 (24%)	3531/14338	_/811	566/3593 (16%)	502/7946 (6%)	1068/12350	
1985	-/1303	1663/5416 (31%)	1671/7815 (21%)	3334/14534	-/718	666/3341 (20%)	612/8891 (7%)	1278/12950	
1990	-/1369	1386/4936 (28%)	1621/7662 (21%)	3007/13967	-/636	658/3242 (20%)	742/8679 (9%)	1400/12557	
1995	-/1143	1225/4795 (26%)	1658/8590 (19%)	2883/14528	-/613	627/3078 (20%)	1178/9737 (12%)	1805/13428	
2000	-/960	915/4198 (22%)	1606/8655 (19%)	2521/13813	-/483	547/2834 (19%)	1255/9588 (13%)	1802/12905	


50-year total* (thousands), mid-1950 to mid-2000: 171 / 1228


-/71	72/252	60/338	132/661	-/42	18/157	21/367	39/566
	(28%)	(18%)			(12%)	(6%)	


 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1950-2000: NEW ZEALAND









NORWAY: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950			•••			•••	•••	•••		
1955	-/1.7	0.4/5.4 (7%)	0.0/7.7 (0.1%)	0.4/15	-/1.1	0.0/4.0	0.0/9.2	0.0/14		
1960	-/1.6	0.7/6.3 (11%)	0.1/9.2 (1%)	0.8/17	-/0.9	0.0/4.1	0.0/10	0.0/15		
1965	-/1.5	0.9/7.4 (12%)	0.3/10 (3%)	1.2/19	-/0.8	0.0/4.2	0.0/11	0.0/16		
1970	-/1.4	1.3/8.1 (16%)	0.6/12 (5%)	1.9/21	-/0.7	0.0/4.3	0.0/12	0.0/17		
1975	-/1.3	1.4/7.9 (18%)	0.9/13 (7%)	2.3/22	-/0.6	0.1/4.1 (2%)	0.0/13	0.1/18		
1980	-/1.1	1.5/8.0 (18%)	1.3/13 (9%)	2.7/23	-/0.5	0.2/3.8 (5%)	0.0/14	0.2/19		
1985	-/1.1	1.6/7.5 (22%)	1.8/15 (12%)	3.4/24	-/0.5	0.3/3.7 (8%)	0.5/16 (3%)	0.8/21		
1990	-/1.1	1.4/6.9 (21%)	1.9/16 (12%)	3.4/24	-/0.5	0.4/3.6 (12%)	0.6/18 (3%)	1.0/22		
1995	-/0.8	1.3/5.9 (23%)	2.2/16 (13%)	3.5/23	-/0.4	0.5/3.3 (16%)	1.0/19 (5%)	1.5/22		
2000	-/0.9	1.2/5.3 (22%)	2.2/15 (14%)	3.4/22	-/0.4	0.6/3.1 (19%)	1.5/19 (8%)	2.1/22		

50-year total* (thousands), mid-1950 to mid-2000: 132 / 1940

-/64	57/343	52/620	108/1027	-/34	9.2/193	15/686	24/913
	(16%)	(8%)			(5%)	(2%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: NORWAY



POLAND: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)			Female (by age)				
Year	0-34	35–69	70+	AII	0–34	35–69	70+	All		
1950			•••			•••		•••		
1955	-/54	6.4/50 (13%)	1.3/32 (4%)	7.7/136	-/40	0.2/41 (0.6%)	0.1/45 (0.2%)	0.3/126		
1960	-/34	9.6/50 (19%)	3.1/32 (10%)	13/117	-/24	0.3/38 (0.9%)	0.2/46 (0.5%)	0.6/107		
1965	-/24	13/58 (23%)	3.2/39 (8%)	16/121	-/15	0.5/41 (1%)	0.2/55 (0.4%)	0.7/111		
1970	-/22	19/67 (28%)	6.6/52 (13%)	25/141	-/13	0.9/42 (2%)	0.5/71 (0.7%)	1.4/126		
1975	-/22	23/72 (31%)	10/64 (16%)	33/158	-/11	1.6/42 (4%)	1.3/85 (2%)	2.9/139		
1980	-/23	31/83 (37%)	14/83 (17%)	45/189	-/11	2.8/44 (6%)	2.1/106 (2%)	4.9/161		
1985	-/20	37/89 (42%)	17/94 (18%)	54/202	-/9.8	3.8/46 (8%)	3.3/124 (3%)	7.1/179		
1990	_/18	45/105 (42%)	15/87 (18%)	60/209	-/7.8	5.1/52 (10%)	4.4/120 (4%)	9.5/179		
1995	-/13	43/105 (41%)	17/87 (20%)	61/206	-/5.9	5.6/49 (11%)	5.2/125 (4%)	11/180		
2000	-/10	37/95 (38%)	20/90 (23%)	57/195	-/4.0	5.9/44 (13%)	5.8/124 (5%)	12/173		


50-year total* (M=millions), mid-1950 to mid-2000: 1.9/ 16M


-/1.3M	1.2/3.7M	0.5/3.2M	1.7/8.2M	-/0.8M	0.1/2.2M	0.1/4.3M	0.2/7.4M
	(33%)	(16%)			(5%)	(2%)	


 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: POLAND









PORTUGAL: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	by age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950						•••		•••		
1955	-/18	0.8/17 (5%)	0.3/16 (2%)	1.1/50	-/15	0.0/12	0.0/22	0.0/49		
1960	-/15	1.1/16 (7%)	0.2/17 (0.9%)	1.2/48	-/12	0.0/11	0.0/23	0.0/47		
1965	-/13	1.5/18 (8%)	0.4/18 (2%)	2.0/49	-/9.5	0.0/12	0.0/25	0.0/46		
1970	-/9.5	2.1/18 (12%)	0.7/20 (3%)	2.8/47	-/7.0	0.0/12	0.0/27	0.0/46		
1975	-/8.0	2.7/21 (13%)	1.2/22 (5%)	3.9/51	-/5.0	0.0/12	0.0/29	0.0/47		
1980	-/6.0	3.1/19 (16%)	1.6/24 (7%)	4.7/50	-/3.3	0.0/11	0.0/31	0.0/45		
1985	-/4.8	3.4/19 (18%)	2.3/27 (9%)	5.7/51	-/2.3	0.0/10	0.0/34	0.0/46		
1990	-/4.2	3.8/19 (20%)	2.8/30 (9%)	6.6/53	-/1.8	0.0/10	0.0/37	0.0/50		
1995	-/4.0	3.9/19 (21%)	3.2/32 (10%)	7.1/54	-/1.5	0.1/9.5 (1%)	0.1/38 (0.2%)	0.2/50		
2000	-/3.2	3.9/18 (22%)	3.7/34 (11%)	7.6/55	-/1.2	0.2/9.0 (2%)	0.3/40 (0.7%)	0.5/50		

50-year total* (thousands), mid-1950 to mid-2000: 199 / 4939

-/476	123/933	74/1152	197/2560	-/335	1.4/561	0.8/1482	2.2/2379
	(13%)	(6%)			(0.2%)	(0.1%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: PORTUGAL



ROMANIA: 1980-2000


Numbers of deaths attributed to smoking / total deaths (thousands)

		Male (k	oy age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950		•••		•••			•••		
1955		•••		•••			•••		
1960		•••		•••			•••		
1965									
1970									
1975									
1980									
1985	-/14	17/53 (31%)	5.4/63 (9%)	22/130	-/8.5	1.6/32 (5%)	0.5/76 (0.7%)	2.2/116	
1990	-/13	20/62 (32%)	4.2/56 (8%)	24/132	-/8.1	2.2/36 (6%)	0.8/72 (1%)	2.9/115	
1995	-/10	24/76 (32%)	5.7/63 (9%)	30/149	-/5.5	2.3/38 (6%)	1.2/79 (1%)	3.5/123	
2000	-/8.7	21/64 (33%)	6.5/64 (10%)	28/136	-/4.7	2.2/34 (7%)	2.6/81 (3%)	4.9/119	

20-year total* (thousands), mid-1980 to mid-2000: 580 / 5183

-/240	412/1288	111/1260	523/2788	-/141	40/699	17/1556	57/2396
	(32%)	(9%)			(6%)	(1%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995



RUSSIAN FEDERATION: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All	
1950		•••					•••		
1955		•••					•••		
1960									
1965									
1970									
1975									
1980	-/129	141/413 (34%)	33/217 (15%)	174/760	-/51	12/248 (5%)	9.0/467 (2%)	21/766	
1985	-/116	162/411 (39%)	43/250 (17%)	206/777	-/49	14/240 (6%)	13/559 (2%)	27/848	
1990	-/102	187/460 (41%)	46/241 (19%)	233/802	-/40	15/251 (6%)	18/563 (3%)	33/854	
1995	-/132	268/753 (36%)	50/283 (18%)	318/1168	-/44	16/347 (5%)	18/645 (3%)	33/1036	
2000	-/124	231/699 (33%)	72/356 (20%)	303/1180	-/39	9.1/311 (3%)	20/695 (3%)	29/1046	

20-year total* (M=millions), mid-1980 to mid-2000: 5.8/ 38M

-/2.5M	4.3/12M	0.9/5.3M	5.2/19M	-/0.9M	0.3/5.9M	0.3/12M	0.6/19M
	(37%)	(18%)			(5%)	(3%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1980-2000: RUSSIAN FEDERATION



SLOVAKIA: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0–34	35–69	70+	All	0–34	35–69	70+	All		
1950						•••	•••			
1955										
1960										
1965										
1970										
1975										
1980										
1985	-/2.4	4.8/12 (39%)	2.6/14 (19%)	7.5/29	-/1.2	0.3/6.1 (4%)	0.4/17 (2%)	0.7/24		
1990	-/2.0	5.9/15 (38%)	1.9/13 (15%)	7.8/30	-/1.0	0.3/7.2 (4%)	0.4/16 (2%)	0.7/24		
1995	-/1.6	5.1/13 (39%)	2.2/13 (17%)	7.4/28	-/0.8	0.2/6.6 (4%)	0.5/17 (3%)	0.7/25		
2000	-/1.4	4.6/13 (35%)	2.6/14 (19%)	7.2/28	-/0.6	0.4/5.9 (6%)	0.5/18 (3%)	0.8/25		

20-year total* (thousands), mid-1980 to mid-2000: 162 / 1051

-/39	100/256	49/272	149/567	-/19	5.1/127	8.5/338	14/484
	(39%)	(18%)			(4%)	(3%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: SLOVAKIA



SPAIN: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)		Female (by age)				
Year	0–34	35–69	70+	All	0–34	35–69	70+	All	
1950		•••	•••			•••		•••	
1955	-/32	5.6/52 (11%)	1.5/52 (3%)	7.1/135	-/25	0.0/40	0.0/68	0.0/134	
1960	-/24	7.6/52 (15%)	3.3/57 (6%)	11/133	-/18	0.3/39	0.0/73	0.3/129	
1965	-/21	8.9/55 (16%)	4.9/62 (8%)	14/138	-/14	0.0/38	0.0/78	0.0/130	
1970	-/17	11/59 (19%)	6.3/68 (9%)	17/144	_/11	0.4/38	0.0/87	0.4/136	
1975	-/16	14/62 (23%)	10/78 (13%)	25/155	-/9.8	0.0/37	0.0/96	0.0/143	
1980	-/12	14/58 (25%)	12/82 (14%)	26/152	-/6.9	0.0/32	0.0/98	0.0/137	
1985	-/10	17/60 (29%)	16/94 (17%)	33/165	-/5.1	0.0/30	0.0/113	0.0/148	
1990	-/13	20/63 (32%)	19/101 (19%)	39/177	-/5.2	0.0/29	0.0/122	0.0/156	
1995	_/11	21/62 (34%)	24/111 (21%)	45/184	-/4.4	0.0/27	0.0/130	0.0/162	
2000	-/8.0	20/59 (33%)	26/123 (21%)	45/189	-/3.3	0.4/25 (2%)	0.0/143	0.4/171	

50-year total* (thousands), mid-1950 to mid-2000: 1236 / 14937

-/901	674/2912	562/3963	1236/7775	-/588	0.0/1723	0.0/4851	0.0/7161
	(23%)	(14%)			(0%)	(0%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: SPAIN



SWEDEN: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (I	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All		
1950			•••	•••						
1955	-/2.8	1.1/13 (8%)	0.4/19 (2%)	1.5/36	-/1.8	0.0/11	0.0/21	0.0/33		
1960	-/2.5	1.6/14 (11%)	0.7/23 (3%)	2.3/40	-/1.6	0.0/10	0.0/24	0.0/36		
1965	-/2.6	1.8/15 (12%)	1.2/24 (5%)	3.0/42	-/1.5	0.0/10	0.0/25	0.0/36		
1970	-/2.4	2.1/16 (13%)	1.9/25 (7%)	4.0/43	-/1.3	0.1/9.7 (0.8%)	0.3/26 (1%)	0.4/37		
1975	-/2.2	2.6/17 (15%)	2.9/29 (10%)	5.5/48	-/1.2	0.3/9.4 (3%)	0.6/29 (2%)	0.9/40		
1980	-/1.9	2.6/16 (16%)	3.4/31 (11%)	6.0/50	-/1.0	0.5/8.8 (6%)	0.8/32 (2%)	1.3/42		
1985	-/1.6	2.6/15 (18%)	3.1/34 (9%)	5.7/50	-/0.9	0.6/7.9 (7%)	0.6/35 (2%)	1.2/44		
1990	-/1.6	2.1/13 (16%)	3.2/34 (9%)	5.3/49	-/0.9	0.7/7.5 (10%)	1.3/38 (3%)	2.0/46		
1995	-/1.2	1.8/11 (15%)	3.3/35 (10%)	5.1/47	-/0.6	0.9/6.8 (14%)	1.9/39 (5%)	2.8/46		
2000	-/1.1	1.5/10 (15%)	3.1/34 (9%)	4.7/46	-/0.5	1.1/6.7 (16%)	2.4/41 (6%)	3.5/48		

50-year total* (thousands), mid-1950 to mid-2000: 257 / 4227

-/104	98/721	109/1408	207/2234	-/61	18/447	31/1485	49/1993
	(14%)	(8%)			(4%)	(2%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: SWEDEN



Year

Year

SWITZERLAND: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (k	oy age)			Female (by age)				
Year	0-34	35–69	70+	All	0–34	35–69	70+	All		
1950				•••		•••		•••		
1955	-/3.0	2.4/11 (22%)	0.7/12 (6%)	3.1/26	-/1.9	0.0/8.1	0.0/15	0.0/25		
1960	-/2.8	2.8/11 (25%)	1.0/13 (8%)	3.8/27	-/1.6	0.0/7.6	0.0/16	0.0/25		
1965	-/2.7	3.2/12 (27%)	1.5/14 (10%)	4.6/29	-/1.6	0.0/7.7	0.0/17	0.0/26		
1970	-/2.7	3.5/12 (29%)	2.1/15 (14%)	5.6/30	-/1.4	0.0/7.2	0.0/19	0.0/27		
1975	-/2.1	3.8/11 (33%)	2.7/16 (17%)	6.4/30	-/1.1	0.1/6.3 (2%)	0.0/19	0.1/26		
1980	-/1.9	3.5/11 (32%)	3.2/18 (18%)	6.6/31	-/0.9	0.2/5.8 (3%)	0.2/22 (0.7%)	0.4/28		
1985	-/1.7	3.3/10 (32%)	3.6/19 (19%)	6.9/31	-/0.8	0.2/5.3 (4%)	0.3/23 (1%)	0.5/29		
1990	-/2.0	3.1/10 (31%)	3.7/20 (18%)	6.8/32	-/0.9	0.3/5.2 (6%)	0.9/25 (3%)	1.2/31		
1995	-/1.7	2.4/9.5 (25%)	3.3/20 (16%)	5.7/32	-/0.8	0.5/5.1 (9%)	0.8/26 (3%)	1.3/32		
2000	-/1.2	2.2/8.8 (26%)	3.3/20 (16%)	5.6/30	-/0.6	0.6/4.9 (12%)	1.2/27 (4%)	1.8/32		

50-year total* (thousands), mid-1950 to mid-2000: 287 / 2848

-/112	151/537	118/821	268/1470	-/62	8.1/325	11/991	19/1378
	(28%)	(14%)			(3%)	(1%)	

 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1955-2000: SWITZERLAND



UKRAINE: 1980-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)			Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	All		
1950				•••						
1955										
1960										
1965										
1970			•••	•••						
1975			•••	•••				•••		
1980			•••	•••						
1985	-/28	51/138 (37%)	18/121 (15%)	69/287	-/13	4.6/86 (5%)	4.8/230 (2%)	9.4/330		
1990	-/26	63/159 (40%)	19/112 (17%)	82/298	-/11	5.6/93 (6%)	8.2/228 (4%)	14/332		
1995	-/32	83/233 (35%)	21/130 (16%)	103/396	-/12	5.5/118 (5%)	7.0/267 (3%)	13/397		
2000	-/28	67/211 (32%)	24/144 (16%)	90/382	-/9.7	3.2/100 (3%)	4.7/266 (2%)	8.0/376		

20-year total* (M=millions), mid-1980 to mid-2000: 1.9/ 14M

-/0.6M	1.3/3.7M	0.4/2.5M	1.7/6.8M	-/0.3M	0.1/2.0M	0.1/5.0M	0.2/7.3M
	(36%)	(15%)			(5%)	(2%)	

^{*}Estimated as 10 times the sum of the annual numbers for 1985 & 1995

1985-2000: UKRAINE



UNITED KINGDOM: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)


		Male (b	y age)		Female (by age)				
Year	0-34	35–69	70+	AII	0-34	35–69	70+	AII	
1950	-/30	47/130 (36%)	11/141 (8%)	58/301	-/24	5.1/97 (5%)	1.7/168 (1%)	6.8/289	
1955	-/23	56/130 (43%)	20/153 (13%)	76/306	-/17	5.9/91 (7%)	2.4/183 (1%)	8.3/290	
1960	-/23	62/135 (46%)	28/151 (19%)	91/309	-/15	7.3/88 (8%)	4.8/191 (3%)	12/295	
1965	-/23	70/146 (48%)	36/155 (23%)	106/323	-/15	11/89 (12%)	6.3/201 (3%)	17/305	
1970	-/20	71/151 (47%)	45/163 (28%)	116/334	-/13	14/90 (15%)	12/218 (6%)	26/321	
1975	-/17	62/142 (44%)	52/176 (30%)	115/335	-/10	16/85 (18%)	15/232 (6%)	31/327	
1980	-/15	54/130 (41%)	55/187 (29%)	109/332	-/9.1	17/79 (21%)	19/241 (8%)	36/329	
1985	-/13	46/117 (39%)	59/202 (29%)	105/332	-/7.5	17/73 (23%)	29/259 (11%)	46/339	
1990	-/13	37/107 (35%)	52/195 (26%)	88/315	-/6.9	16/68 (24%)	32/252 (13%)	49/327	
1995	-/11	28/93 (30%)	48/206 (24%)	76/311	-/6.0	13/60 (22%)	39/269 (15%)	53/335	
2000	-/10	21/83 (25%)	42/197 (21%)	63/290	-/5.3	11/54 (21%)	40/259 (15%)	51/318	


50-year total* (M=millions), mid-1950 to mid-2000: 6.3/ 32M


-/0.9M	2.6/6.3M	2.2/8.9M	4.8/16M	-/0.6M	0.6/4.0M	0.9/11M	1.5/16M
	(42%)	(24%)			(16%)	(8%)	


 $^{^{\}star}\textsc{Estimated}$ as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1950-2000: UNITED KINGDOM









UNITED STATES: 1950-2000

Numbers of deaths attributed to smoking / total deaths (thousands)

		Male (I	oy age)		Female (by age)				
Year	0-34	35–69	70+	All	0-34	35–69	70+	AII	
1950	-/122	62/398 (16%)	8.2/308 (3%)	70/828	-/85	0.0/248	0.0/292	0.0/625	
1955	-/119	83/403 (20%)	17/350 (5%)	99/873	-/79	0.7/241 (0.3%)	0.0/336	0.7/656	
1960	-/123	109/439 (25%)	30/413 (7%)	139/976	-/81	3.6/254 (1%)	0.0/401	3.6/736	
1965	-/118	131/460 (28%)	47/458 (10%)	178/1035	-/74	12/265 (5%)	0.8/454 (0.2%)	13/793	
1970	-/119	153/482 (32%)	66/477 (14%)	219/1078	-/68	27/276 (10%)	6.5/498 (1%)	34/843	
1975	-/109	155/457 (34%)	83/485 (17%)	237/1051	-/56	40/262 (15%)	15/524 (3%)	55/842	
1980	-/110	155/439 (35%)	104/526 (20%)	259/1075	-/54	54/261 (21%)	37/600 (6%)	91/915	
1985	-/98	152/425 (36%)	122/575 (21%)	274/1098	-/48	66/261 (25%)	65/680 (10%)	132/989	
1990	-/103	147/415 (35%)	133/595 (22%)	280/1113	-/48	72/257 (28%)	102/730 (14%)	174/1035	
1995	-/93	136/423 (32%)	145/657 (22%)	281/1173	-/44	74/262 (28%)	140/833 (17%)	215/1139	
2000	-/74	118/405 (29%)	150/699 (22%)	269/1178	-/38	73/268 (27%)	170/920 (18%)	243/1226	

50-year total* (M=millions), mid-1950 to mid-2000: 15/96M

1950-2000 by age & sex:

-/5.4M 6.6/22M 4.1/25M 11/52M -/3.0M 1.9/13M 2.2/28M 4.1/44M (30%) (16%) (15%) (8%)

^{*}Estimated as 10 times the sum of the annual numbers for 1955, 1965, 1975, 1985 & 1995

1950-2000: UNITED STATES

